
James Webb Space Telescope





Tom Donaldson, Jason Kalirai, Jeff Valenti




IVOA Northern Spring Interop, Cape Town

#JWST



The Observatory






Mirror Revealed



Lead project scientist John Mather admires all 18 segments of the 6.5m mirror


Orbit


- Orbit is around the 2nd
 Lagrange point (L2)
- About 1.5 million km from earth



Field of Regard

- ~40% of the sky at any one time
- Every point on the sky is visible for at least 100 days per year.


















Field of View



Imaging Modes

Mode	Instrument	Wavelength (micron)	Pixel Scale (arcsec)	Field of View
Imaging	NIRCam	0.6-2.3	0.032	2.2×4.4
		2.4-5.0	0.065	2.2×4.4
	NIRISS	0.9-5.0	0.065	2.2×2.2'
	MIRI	5.0-28	0.11	1.2×1.9'
Coronagraphy	NIRCam	0.6-2.3	0.032	20×20"
		2.4-5.0	0.065	20×20"
	MIRI	10.65	0.11	4Q: 24x24"
		11.4	0.11	4Q: 24x24"
		15.5	0.11	4Q: 24x24"
		23	0.11	Lyot: 30×30"
Aperture Mask Interferometry	NIRISS	3.8-4.8	0.065	2.2×2.2'

Spectroscopic Modes

Mode	Instrument	Wavelength (micron)	Resolution (R= λ / Δ λ)	Field of View
Single Slit Spectroscopy	NIRSpec	0.6-5.0	100, 1000, 2700	0.4x3.8" 0.2x3.3" 1.6x1.6"
	MIRI	5.0-12.0	100	0.6×5.5" slit
Multi-Object Spectroscopy	NIRSpec	0.6-5.0	100, 1000, 2700	3.4x3.4' 0.2x0.5" shutters
Slitless Spectroscopy	NIRISS	1.0-2.5	150	2.2×2.2'
		0.6-2.5	700	single object
	NIRCam	2.4-5.0	1700	2.2×2.2'
Integral Field Unit Spectroscopy	NIRSpec	0.6-5.0	100,1000,2700	3.0×3.0"
	MIRI	5.0-7.7	3500	3.0×3.9"
		7.7-11.9	2800	3.5×4.4"
		11.9-18.3	2700	5.2×6.2"
		18.3-28.8	2200	6.7×7.7"


Multi-Object Spectroscopy in Crowded Environments








Micro-Shutter Assembly (MSA) has 248,000 microshutters!





Multi-Object Spectroscopy in Crowded Environments



on the sky

+ Targets in Operable Shutters x Targets Outside Shutters

JWST Products and Services


- JWST data processing and archiving done at STScI
- A variety of products and services will be provided:
 - Images
 - Spectra
 - Catalogs
 - Web-based user interfaces
 - Data Analysis Tools and Software
 - Mission documentation

Image Products

- Multiple levels of image products are created during processing.
 Publicly accessible data begins at level 1b.
 - 1a: Original FITS files
 - 1b: Uncalibrated FITS files
 - 2a: Ramp calibrated exposure FITS files
 - 2b: Fully calibrated exposure FITS files
 - 3: Calibrated combined (dithers, mosaics) FITS files
 - VO metadata clearly defined for combined products?
 - 4: Products resulting from additional analysis
- For each data product, a variety of calibration files and other artifacts must be publicly available to support custom processing.

Image Cubes

Up-the-ramp data collection results in 4D cubes



Swade, et al., <u>2012ASPC..461..225S</u>

Time Series Images for Transits

- 2 day long exposure product may be stored in a single 35GB FITS file with time-tags as extension.
- Levels 1-3 data products would total 180GB
- Most transit observations will be shorter and have smaller products.

Spectral Data Products


- 1D extracted per exposure/integration/NIRSpec background
- 2D for non-IFU spectra
- 3D for IFU data
- Combined level-3 1D and 2D spectra
 - MSA observations have complex dither strategies
- Separate MSA spectral product for each source

Catalogs

- Source catalogs extracted from each level-3 image
 - NIRCam catalogs needed for MSA observation planning
- JWST Source Catalog will be produced by combining catalogs from individual images.

User Interfaces


Primary UI is the MAST Discovery Portal (mast.stsci.edu)



Data Analysis Tools and Software

- Processing pipeline software
- New interactive analysis tools
- Astropy features
- Server-side analysis (bringing code to the data)

Data Analysis Tool Example: Cube Viewer



Products and Services Summary

- Images
- Spectra
- Catalogs
- Web-based user interfaces
- Data Analysis Tools and Software
- Mission documentation

#JWST